

Tuberculosis 95 (2015) 246-250

Contents lists available at ScienceDirect

## Tuberculosis

journal homepage: <http://intl.elsevierhealth.com/journals/tube>

### MOLECULAR ASPECTS

#### MicroRNA-155 knockout mice are susceptible to Mycobacterium tuberculosis infection

Hiroki Iwai <sup>a</sup>, Keiji Funatogawa <sup>b</sup>, Kazunori Matsumura <sup>a</sup>, Masako Kato-Miyazawa <sup>a</sup>,  
Fumiko Kirikae <sup>a</sup>, Kotaro Kiga <sup>d</sup>, Chihiro Sasakawa <sup>c, d, e</sup>, Tohru Miyoshi-Akiyama <sup>a</sup>,  
Teruo Kirikae <sup>a, \*</sup>

<sup>a</sup> Department of Infectious Diseases, Research Institute, National Center for Global Health and Medicine, Toyama, Shinjuku, Tokyo 162-8655, Japan

<sup>b</sup> Tochigi Prefectural Institute of Public Health, Tochigi 329-1196, Japan

<sup>c</sup> Nippon Institute for Biological Science, Tokyo 198-0024, Japan

<sup>d</sup> Division of Bacterial Infection, The Institute of Medical Science, The University of Tokyo, Tokyo 108-8639, Japan

<sup>e</sup> Medical Mycology Research Center, Chiba University, Chiba 260-8673, Japan

### article info

Article history: Received 1 December 2014 Accepted 15 March 2015

Keywords: Mycobacterium tuberculosis microRNA miR-155

### summary

MicroRNAs (miRNAs) are short, conserved, non-coding RNA molecules that repress translation, followed by the decay of miRNA-targeted mRNAs that encode molecules involved in cell differentiation, development, immunity and apoptosis. At least six miRNAs, including microRNA-155 (miR-155), were upregulated when bone marrow-derived macrophages from C57BL/6 mice were infected with Mycobacterium tuberculosis Erdman. C57BL/6 mice intravenously infected with Erdman showed up-regulation of miR-155 in livers and lungs. Following infection, miR-155-deficient C57BL/6 mice died significantly earlier and had significantly higher numbers of CFU in lungs than wild-type mice. Moreover, fewer CD4<sup>+</sup> T cells, but higher numbers of monocytes and neutrophils, were present in the lungs of Erdman-infected miR-155 knockout (miR-155<sup>-/-</sup>) than of wild-type mice. These findings indicated that miR-155 plays a critical role in immune responses to M. tuberculosis.

© 2015 Elsevier Ltd. All rights reserved.

\* Corresponding author. Tel.: +81 3 3202 7181; fax: +81 3 3202 7364. Th17 and regulatory T cells [13]. miR-155 regulates a wide spectrum

E-mail address: [tkirikae@ri.ncgm.go.jp](mailto:tkirikae@ri.ncgm.go.jp) (T. Kirikae).

<http://dx.doi.org/10.1016/j.tube.2015.03.006> 1472-9792/© 2015 Elsevier Ltd. All rights reserved.